

Phenomenological Theory of Survival

Mark Ya. Azbel'

School of Physics and Astronomy, Tel Aviv University,

Ramat Aviv, 69978 Tel Aviv, Israel

(February 2, 2008)

Abstract

Theoretical analysis proves that human survivability is dominated by an unusual physical, rather than biological, mechanism, which yields an exact law. The law agrees with all experimental data, but, contrary to existing theories, it is the same for an entire species, i.e., it is independent of the population, its phenotypes, environment and history. The law implies that the survivability changes with environment via phase transitions, which are simultaneous for all generations. They allow for a rapid (within few percent of the life span) and significant increase in the life expectancy even above its value at a much earlier age.

Mortality is one of the most universal and important phenomena in biology. Human mortality is extensively studied over two centuries [1–5] and is arguably the best statistically quantified biological phenomenon. There exist several evolutionary theories of aging [6] (the first one [7] is only 50 years old). Yet, C. Franceschi (in ref. [8]) notes: "Longevity is a trait with some peculiarities because of the unnecessary nature of aging". This paper proves that, in contrast to all existing theories, human mortality is dominated by an unusual physical (rather than biological) mechanism. Presumably, the same mechanism dominates mortality of laboratory fly populations.

I study a large and diverse amount of over 150,000 data points from 1528 period life tables, for 3 races on 4 continents in 16 countries during over a century of their history, 36 cases total (here and on "a case" denotes the population of a given sex in a given country at any time in its history; in the USA white, black and total populations are considered as separate cases). Each period life table presents in any given case the probability l_x to survive to any (yearly) age x in a given calendar year. To escape any arbitrary scaling or adjustment, I use life table variables only.

The survivability l_x depends on age x , calendar year t and a complete set A of all parameters, comprehensively describing a considered population, i.e. its phenotypes and their heterogeneous social, medical, dietary, etc. conditions during the previous x years (from $t - x$ till t). So, $l_x = l_x(t, A)$. Mortality rate, and thus its statistics and accuracy, are low in early age. The number of survivors, and thus accuracy, decrease in old age. So, middle age survivability, e.g., l_{40} , is determined more accurately. Present $l_x(t, A)$ and $l_{40}(t, A)$ from the same life table as the ordinate and abscissa of a point. According to Fig. 1, for different races, countries, continents, histories l_x is dominated by the function $l_x(l_{40})$, which depends only on l_{40} , i.e. invariant to A and is therefore the same for an entire species. The deviation of l_x from this function depends on A (and thus on population phenotypes, living conditions, history). It is relatively small. From now on I denote $l_x(l_{40})$ as "the invariant survivability" and, unless specifically stated otherwise, consider this (dominant) fraction of survivability only.

By Fig. 1, at any age and in every case, l_{40} significantly changes with living conditions. Thus, since living conditions of different groups in a given case are different, these groups may have different probabilities l'_x to survive to x years (the value of l_x in the population is the average l'_x ; $0 \leq l_x, l'_x \leq 1$). Suppose the probability density of a given l'_{40} in the population is $c(l_{40}, l'_{40})$, i.e.

$$\int_0^1 c(l_{40}, l'_{40}) dl'_{40} = 1, \quad l_x(l_{40}) = \int_0^1 l'_x \cdot c(l_{40}, l'_{40}) dl'_{40} \quad (1)$$

By Eq. 1, $0 \leq \min(l'_{40}) \leq l_{40} \leq \max l'_{40}$. So, $l'_{40} = 0$ when $l_{40} = 0$ and $l'_{40} = 1$ when $l_{40} = 1$. Thus, invariance, which allows one to introduce c and which yields Eq. 1, implies that any population is homogeneous in l'_{40} at the distribution boundaries. In virtue of invariance, the dependence of l'_x on l'_{40} is the same as the dependence $l_x(l_{40})$, i.e., $l'_x = l_x(l'_{40})$. This, by Eq. 1, yields a remarkable symmetry of the invariant survivability to the transformations, specified by the function c of two variables:

$$l_x \left[\int_0^1 l'_{40} \cdot c(l_{40}, l'_{40}) dl'_{40} \right] = \int_0^1 l_x(l'_{40}) \cdot c(l_{40}, l'_{40}) dl'_{40} \quad (2)$$

Such invariance is sufficiently restrictive to mathematically accurately determine $l_x(l_{40})$ (as an invariant of survivability dynamics with respect to any population and environmental changes). Indeed, in a general case the function $l_x(l_{40})$, which yields Eq. 2, must be linear. A special case of $c(l_{40}, l'_{40}) = c' \cdot \delta(l'_{40} - l') + c'' \cdot \delta(l'_{40} - l'')$ yields $c' + c'' = 1$ by Eq.1, and $l_x(l_{40}) = l_x(c' \cdot l' + c'' \cdot l'') = c' \cdot l_x(l') + c'' \cdot l_x(l'')$ by Eq. 2. The latter equation implies $dl_x/dl_{40} = dl_x/dl'_x = \text{const.}$ On the other hand, a linear $l_x(l_{40})$ yields Eq. 2 with any arbitrary c in virtue of Eq. 1. Similarly, if an entire population is distributed in a certain interval of l_{40} , then $l_x(l_{40})$ is linear in this interval, and the population is homogeneous at its ends (as it was in the previous case). Correspondingly, if any population is homogeneous at certain l_{40} points only, then an entire population is distributed within one of the intervals $\lambda_s < l_{40} < \lambda_{s+1}$ ($s = 0, 1$ is its ordinal number):

$$\int_{\lambda_s}^{\lambda_{s+1}} c(l_{40}, l'_{40}) dl'_{40} = 1; \quad l_{40} = \int_{\lambda_s}^{\lambda_{s+1}} l'_{40} \cdot c(l_{40}, l'_{40}) dl'_{40} \quad (3)$$

and $l_x(l_{40})$ is linear within these intervals. When $l_{40} = \lambda_{s+1}$, an entire population of all ages moves to the next interval. Clearly, such piecewise $l_x(l_{40})$ satisfies Eq.2. When $\lambda_0 = 0, \lambda_1 = 1$, this case reduces to the previous one. If an infinite number of edge points in Eq.3 condenses and forms a continuous interval (where any population is homogeneous), then the invariant $l_x(l_{40})$ in this interval remains undetermined. Of course, the possibility to distinguish this case from the case of a large number of short linear segments is limited by the disregarded non-invariant contributions to survivability, but demographic studies prove that survivability is heterogeneous and exclude the former case (of a population which is homogeneous in a finite interval). Thus, in a general case

$$l_x = R_x^{(s)} l_{40} + a_x^{(s)}, \quad \lambda_s \leq l_{40} \leq \lambda_{s+1}. \quad (4)$$

So, except for the number of linear segments, the very existence of the invariant survivability allows one to establish its exact law, without any experiments, approximations, and assumptions.

Equation (4) is the implication of invariance only. Such invariance was also demonstrated [9] for medfly and fruitfly families whose different populations were extensively studied in different conditions. Fly statistics is rather low, thus their l_x was studied as a function of the life expectancy at birth (which averages l_x over different ages and is therefore more smooth). The study was based on fly populations hatched the same day. However, since in most cases fruitflies were kept in stationary conditions, it suggests that the invariant survivability law is general for humans and flies.

Piecewise linear law (4) and its age independent invariant intersections are the main predictions of this paper. They are verified with all experimental data - see, e.g., l_1, l_{60}, l_{80} vs l_{40} in Fig.2. Piecewise linear law agrees with (but has never been suggested in) demographic approximations [3]. Slope jumps in Eq.(4) and Fig.2 are consistent with ref [5]. However, in contrast to the qualitative observations in ref. [5], accurate Eq.(4) yield quantitative predictions. Since all l_x belong to the same calendar year, their slope jumps are simultaneous for all generations. Indeed, although the intersections in Fig.2 are reached at different times

in different countries for different sexes, they are simultaneous (to the invariance accuracy) at different ages, and (in l_{40}) for different populations. The invariant law allows for the prediction of intersections. For instance, the extrapolation of l_{80} beyond $\lambda_1 = 0.95$ in Fig. 1 yields $l_{40} = 1$ (while l_{80} is finite), i.e. implies no deceased until 40y. This is hardly possible, and suggests a crossover, which is indeed seen in Fig.2 at $\lambda_2 = 0.97$. The extrapolations of l_{60} and l_{80} to $l_{40} < 0.94$ yield $l_{60} = 0$ and $l_{80} = 0$, i.e., no survivors beyond 60y (when $l_{40} = 0.27$) and 80y (when $l_{40} = 0.42$). This is never true, and suggests an intersection at $0.4 < l_{40} < 0.7$.

To elucidate the nature of the invariant law, present Eq. (4) in a different form:

$$l_x = c_x l_x^{(s)} + (1 - c_s) l_x^{(s+1)} \quad \text{when} \quad l_x^{(s)} \leq l_x \leq l_x^{(s+1)} \quad (5)$$

Here $l_{40}^{(s)} = \lambda_s$; $l_x^{(s)} = l_x(\lambda_s)$ from Eq.(4); c_s reduces to $l_x^{(s)}$ and l_{40} [thus, by Eq. (3), to c]. Equation (5) accurately separates "nature" and "nurture" in survival. "Nature" reduces to the fixed set of the intersection survivabilities. The set depends only on age and is invariant, i.e., independent of phenotypes (and thus of a specific DNA sequence at least in an entire species), their living conditions and life history. The dependence of the set on age is not determined by invariance, but invariance implies that it is the same for at least an entire species. (Moreover, it scales onto the same functions for species as remote as humans and flies [9]). Thus, it does not change at least as long as the species does not evolve into a different species (demographic data in Fig.2 verify it for 100-150 years). So, the set must be inheritable. Such set reminds of the body temperature (which in any living conditions is the same with few percent accuracy) of an entire class of birds and a subclass of placental mammals. Presumably, both the set and the body temperature are genetically determined, and independent of a specific DNA sequence. But the set, unlike the body temperature, strongly depends on age. "Nurture" distributes the survivability l_x at a given age between two adjacent intersection survivabilities. The concentration $c_s (0 \leq c_s \leq 1)$ is age independent (and may be related to, e.g., l_{40}). Thus, survivability follows environment (in particular, a new intersection survivability emerges) simultaneously for all generations.

Equation (5) relates c_s , and thus l_x , to l_1 . Since l_x is the survival probability in the same calendar year as l_1 , so, by Eq. (5), the survivability accurately and rapidly follows the change in environment according to the value of l_1 (i.e., the infant mortality $q_0 = 1 - l_1$), which is established in less than two years. Indeed, whatever the difference in environmental factors is, close values of infant mortality imply very close fractions of deceased at any given age in 1885 Swedish and 1947 Japanese females, despite of their different races, continents, countries, and 62 year gap in their different history. Since l_1 depends on the environment in a given year only, the invariant survivability l_x is reversible (an entire survival curve comes back when q_0 changes non-monotonically and returns to its previous value), and statistically independent of the life history during its x years, despite country specific, highly and irregularly changing, non-monotonic and non-stationary living conditions. World wars and epidemics, e.g., flu in 1918 Europe, significantly decrease l_x . (For instance, in 1915 the probability for a French male to survive to 80y was 5 times less than in 1913, twice less than in 1917, and 3.5 times less than for a 1861 Swedish male). Yet, they just slightly shift the plots (mostly vertically, and relatively little) in Fig.2. In a couple of years (which estimate the relaxation time at few percents of the life span) all plots restore their invariant dependence, i.e., the memory of the previous life history is erased. Accurate reversibility of the invariant survivability does not decrease with aging, even in old age. Unless such reversibility is related to some perfect biological rehabilitation (which is hardly possible), it implies an adiabatic change in a certain thermodynamic equilibrium. This is consistent with its relaxation time (rapid compared to the life span, but enormous on a microscopic scale). Equation (5) reduces l_x to the fixed set of $l_x^{(s)}$. Since $l_x^{(s)}$ reversibly change into each other, they are related to different equilibrium thermodynamic states of the same system, i.e. to different phases. Thus, Eq. (5) relates the invariant survivability to a certain phase equilibrium, and the jumps in Fig.1 slopes to the emergence of a new phase. The phase concentration c_s , which Eq.(5) reduces to, e.g., l_1 , is independent of age. This suggests that the age dependence of a $l_x^{(s)}$ is related to the difference in survivabilities provided by the same phase at different ages. Quasi-equilibrium phases in a living homeostatic being might

be related to a meso- (e.g., a cell) or microscopic (e.g., DNA configuration) scale.

While the existence of survivability phases is accurately proven, their microscopic nature, as well as that of the parameter which determines their concentrations, remains unknown (as "units of heredity" were to Mendel). However, any phase equilibrium may be reversibly manipulated. This means that human life expectancy may be rapidly (within few years) reversed to its value at a much earlier age. (Note that in just 8 years from 1947 till 1955 the life expectancy of Japanese females increased 26% at birth, 15% at 60y, and 20% at 80y).

Presumably, the change in phase concentrations may affect other than survivability characteristics, in particular, aging and disease resistance. Indeed, Eq.(5) is not violated even in Japan prior to 1949 (see Fig.1) and in Finland from 1890 till 1940 (see Fig.2), although their mortality has a strongly tubercular age pattern during this period.

The suggested phase equilibrium nature of survivability implies that non-equilibrium (in particular, sufficiently non-stationary and heterogeneous conditions which depend on age, sex, social mobility, immigration and other factors), may lead to more than two adjacent phases and to hysteresis in the adjustment. The contribution of non-equilibrium phases is most pronounced in old age, when the difference in $l_x^{(s)}$ is significantly higher. This agrees with Fig.2. Hysteresis yields small jumps in l_x and shifts of the intersections, which are indeed present in certain cases. The accuracy of the invariant law estimates maximal concentrations of "extra" phases. They are lower for females, presumably suggesting lower heterogeneity of female populations.

Survivability independence of life history implies no correlation between early and old age invariant mortalities in a given cohort (born the same year). This disagrees with evolutionary theories of aging [6]. Stochastic mutation accumulation [7] theory is inconsistent with reversibility and rapid accurate survivability change with environment. Optimal allocation of metabolic resources (pleiotropy and disposable soma theories) implies strong correlation between survival in young and old age in populations born the same year. This is inconsistent with the survivability independence of life history. Reversibility is also inconsistent with mortality theories (telomeres, oxygen consumption, free radicals, somatic mutations). Al-

though natural mortality in the wild is mostly due to extrinsic hazards, invariant mortality, which dominates in species as remote as humans and flies (in certain protected environment), and which rapidly, accurately and simultaneously in all generations changes with environment, calls for biological, evolutionary and microscopic physical theories.

These conclusions, as well as an unusual mechanism of mortality, are accurate implications of the exact law, which agrees with all demographic approximations and studies [3,5]. For instance, the infant mortality is widely appreciated by demographers as a sensitive barometer of environmental conditions. However, demographic approximations are developed primarily as a useful tool of maximally accurate estimation and forecast. They are often country and time specific, and approximate specific demographic data better than Eq. (4), but they do not consider nor care about the underlying general law and its mechanism. In contrast, my goal is the exact law, albeit of the invariant mortality only (to discover inertia, one must disregard friction!) and its mechanism, which yield new biological insights.

Acknowledgement

This paper was stimulated by enlightening discussions with Profs. Y. Aharonov, S.M. Jazwinski, S. Horiuchi, D.S. Thaler, A. Libchaber, and B. Tsirelson. I am grateful to Prof. S. Horiuchi, S. Smallwood, A. Hanika, K. Anderson, S. Helgadottir, C. Capocci, E. Waeytens, N. White, U. Wegmueller, V. Kannisto, M. Nieminen, K. Deaves for life tables; to I. Kolodnaya for crucial technical assistance; A. Kobi for computer simulations, J. & R. Meyerhoff chair for financial support.

REFERENCES

- [1] Halley,E., Phil. Trans. Roy. Soc.**17**, 596 (1693); Euler, L., Histoire de l'Academie Royale des Sciences et Belle- Lettres, 144-164 (1760).
- [2] Gompertz, B., Phil. Trans. Roy. Soc. **A115**, 513 (1825)
- [3] Coale, A.J., Demeny, P., Vaughan, B., Regional Model Life Tables and Stable Populations, 2nd ed. (academic, NY, 1993); Lee, R.D., Carter, L.R., J. Am. Stat. Assoc. **87**, 659 (1993); Indirect Techniques for Demographic Estimation, Manual X, pp.12-21, UN publication (1993), and refs. therein.
- [4] Australian life tables (1999), Australian Government Actuary, Canberra; Tables de Mortalité (1999), Statistics Belgium, Brussels; Life tables for Japan (1994), Ministry of Health and Welfare, Tokyo; Berkeley Mortality Database, <http://demog.berkeley.edu/wilmoth/mortality>
- [5] Kannisto, V. (1994), Development of oldest-old mortality, 1950-1990, Odense, Denmark (Odense Univ. Press). Horiuchi, S., Wilmoth, J.R. (1998). Demography **35** 4, 391-412. Wilmoth, J.R., Horiuchi, S. (1999), Demography **36** (4), 475-495. Tuljapukar, S., Li, N., Boe, C. , Nature **405**, 789-792 (2000). Horiuchi, S. in Health and Mortality: Issues of Global Concern, 54 (1999), UN, NY, and refs. therein.
- [6] Kirkwood, T.B.L., Austad, S.N. , Nature , 233-238 (2000).
- [7] Medawar, P.B., An Unsolved Problem of Biology, Lewis, London, (1952).
- [8] Koenig, R. Science 291, 2074 (2001).
- [9] Azbel, M. Ya., PNAS USA **96** 3303, 15368 (1999)

FIGURES

FIG. 1. Probabilities (vertical axis, upper, middle and lower curves correspondingly) to survive to 1, 60, 80 years vs l_{40} (horizontal axis) for females in 1909-1997 Australia, 1880-1998 Belgium, 1891-1996 Japan, 1861-1995 Sweden. Swedish and Japanese females are denoted by full square, open square, and full triangle, open triangle; others by full circle, open circle; open signs denote 1914-1919; 1939-1947. Solid lines are linear regressions, which verify predicted piecewise linearity.

FIG. 2. Same as in Fig. 1, but for all cases (males and females in 1909-1997 Australia, 1880-1998 Austria; 1880-1998 Belgium, 1950-1987 Canada and provinces; 1851-1998 England and Wales; 1881-1998 Finland, 1898-1995 France; 1871-1994 Germany; 1841-1998 Iceland; 1925-1992 Ireland; 1891-1996 Japan, 1846-1998 Norway; 1930-1992 Scotland; 1861-1995 Sweden, 1878-1993 Switzerland; 1900-1997 USA, white, black, and total population.) To amplify invariance and piecewise linear dependence, some of the linear segments are slightly rotated and shifted. (This does not violate piecewise linearity.)





